

HD-EXTRA BIO

detergent sanitizer



- CLINIMAX HD-EXTRA BIO is the new heavy duty detergent sanitizer, designed to offer the user the peace of mind and cost efficiency associated with all CLINIMAX quality products.
- CLINIMAX HD-EXTRA breaks down surface tension, weakens bacteria cell walls, allowing the optimum antimicrobial and disinfectant action of its active Biocide to take place.
- Applied either manually or by pressure washer, CLINIMAX HD-EXTRA BIO can reduce water consumption and provide quick and efficient disinfection.
- User and environmentally friendly, CLINIMAX HD-EXTRA BIO is of very low toxicity at the recommended dilution applications.

APPLICATIONS

For the simple, safe and effective cleaning of all surfaces prior to Disinfection. For Cleaning and Sanitizing farm equipment, vehicles, foot and wheel dips, hatcheries, dairies, farms, stables, catteries, kennels, other animal accommodation, abattoirs and food processing plant. In a disease prevention program, use CLINIMAX HD-EXTRA BIO as a cleaner and sanitizer prior to disinfection. Use as part of your regular cleaning routine.

Available in 5L and 25L re-sealable containers.

INSTRUCTIONS FOR USE:

GENERAL USE AND PRESSURE WASHING. For general use (floors, walls, etc) use CLINIMAX HD-EXTRA BIO at a dilution rate of **1:100** (100ml in 10 litres of water).

PRE-DISINFECTION CLEANING & NORMAL SURFACE WASHING. For cleaning and sanitizing all farm equipment use CLINIMAX HD-EXTRA BIO at a dilution rate of **1:200 to 1:100** (50-100ml per 10 litres of water.)

FINAL STAGE CLEANING & CLEANING HEAVILY SOILED SURFACES. For final stage clean out of animal housing, for cleaning and sanitizing abattoir equipment and surfaces, and for cleaning all heavily soiled surfaces, buildings and equipment, use CLINIMAX HD-EXTRA BIO at a dilution rate of **1:100 to 1:50** (100-200ml per 10 litres of water), depending on the degree of soiling. Apply CLINIMAX HD-EXTRA BIO to the soiled surfaces, leave to soak and penetrate the soiling, then remove by pressure washing.

CLEANING EXTREMELY SOILED SURFACES. For removing extreme soiling from surfaces, grease and oil use CLINIMAX HD-EXTRA BIO at a dilution rate of between **1:50 and 1:1**. Apply to the soiled surface, then wash with hot water. Scrub with a stiff brush to facilitate emulsification and oil removal, wash off with water and repeat if necessary.

DEFRA APPROVALS

This disinfectant has been approved for use in England by the Secretary of State for the Department for Environment, Food and Rural Affairs and in Scotland by the Scottish Ministers for the purposes of:

General orders made under the Animal Health Act 1981 at a dilution rate of one part of this preparation to 100 parts of water.

The contents of this container are guaranteed to be of the same quality and composition as the sample submitted for approval testing.

VICTOR®

broad spectrum disinfectant

&

HD-EXTRA BIO

detergent sanitizer



DOES NOT CONTAIN PROHIBITED SUBSTANCES UNDER THE JOCKEY CLUB RULES OF RACING

A complete
cost effective
disinfection and
disease control
program from



For further information contact
www.clinimax.net

CLINIMAX
health and hygiene technology

successful science through innovation and discovery

CLINIMAX
health and hygiene technology

successful science through innovation and discovery

VICTOR[®]

broad spectrum disinfectant



- **VICTOR** is a broad spectrum disinfectant, designed to offer the user the peace of mind and cost efficiency associated with all CLINIMAX quality products.
- **VICTOR** is one of the most powerful disinfectants ever developed.
- **VICTOR** can be used to clean and disinfect surfaces in a simple one step operation.
- User and environmentally friendly, **VICTOR** is of very low toxicity at the recommended dilution applications.

APPLICATIONS

For the simple, safe and effective control of viruses, bacteria, fungi, yeasts and moulds in **farms, veterinary surgeries, operating theatres, foot and wheel dips, hatcheries, dairies, stables, catteries, kennels** and **other animal accommodation**, for the cleaning and disinfection of all surfaces, equipment and utensils.

Available in 1kg, 5kg and 15kg re-sealable packs.

DEFRA APPROVALS

This disinfectant has been approved for use in England by the Secretary of State for the Department for Environment, Food and Rural Affairs and in Scotland by the Scottish Ministers for the purposes of the Foot and Mouth Disease Order 1983 at the dilution rate of one part of this preparation to 1300 parts of water;

the Swine Vesicular Disease Order 1972 at the dilution rate of one part of this preparation to 200 parts of water;

the Diseases of Poultry (England) Order 2003 and the Diseases of Poultry (Scotland) Order 2003. The Avian Influenza and Influenza of Avian Origin in Mammals (England) Order 2006 and the Avian Influenza and Influenza of Avian Origin in Mammals (Scotland) Order at a dilution rate of one part of this preparation to 280 parts of water;

General orders made under the Animal Health Act 1981 at a dilution rate of one part of this preparation to 120 parts of water.

The contents of this container are guaranteed to be of the same quality and composition as the sample submitted for approval testing.

This preparation is also approved by the Department of Agriculture & Rural Development, Northern Ireland as a disinfectant under the Diseases of Animals (Northern Ireland) Order 1981 at the dilution rate stated above.

INSTRUCTIONS FOR USE:

FINAL STAGE DISINFECTION: Apply VICTOR at a dilution rate of **1:100 (1%)** using a pressure washer (at 300 psi) or other spraying equipment at an application rate of 300ml/m².

CLEANING & DISINFECTING BUILDINGS: As part of specific control programs, use a pressure washer (at 300 psi) or other spraying equipment at a dilution rate of **1:200 (0.5%)** at an application rate of 500ml/m².

CLEANING & DISINFECTING EQUIPMENT: For equipment on farms, hatcheries, food processing plants etc, use a pressure washer (at 300 psi) or other spraying equipment at a dilution rate of **between 1:100 (1%) and 1:200 (0.5%)** depending on soiling levels at an application rate of 300ml/m².

AERIAL DISINFECTION: In cases where there is a risk of cross infection, particularly during outbreaks of respiratory disease, use either a pressure washer at low pressure, a ULV sprayer or a knapsack sprayer at a dilution rate of **1:200 (0.5%)** to fill the atmosphere with a fine mist at an application rate of 1L of solution per 100m³ of air space.

FOGGING IN PLACE OF FORMALIN FUMIGATION:

A thermal fogging machine should be used to apply Victor at a dilution rate of **1:25 (4%)** to 25ml/m² of floor space.

WATER SYSTEM SANITATION:

For final stage cleanout, soak water systems for a minimum of 1 hour in a solution of **1:200 (0.5%)** and then flush out with clean water. For continuous sanitation, apply VICTOR through a dosing system at a dilution rate of **1:1000 (0.1%)**

FOOTDIPS: Use at a dilution rate of **1:100** and replenish once heavy soiling occurs. Discard after a maximum of 1 weeks use.

This product has been approved by the Minister for Agriculture & Food, Dublin, as a disinfectant for the purposes of the Diseases of Animals Act, 1966 and Orders made thereunder.

Approved Uses and Dilution Rates

Foot and Mouth Disease Orders in the proportion of one part of this preparation to 1300 parts of water.

Swine Vesicular Disease Orders in the proportion of one part of this preparation to 200 parts of water.

Diseases of Poultry Orders in the proportion of one part of this preparation to 280 parts of water.

General Orders made under the Animal Health Act 1966 in the proportion of one part of this preparation to 120 parts of water.

The contents of this container are guaranteed to be of the same quality and composition as approved by the Minister for Agriculture & Food, Dublin.

PLEASE NOTE

Dilution rates for disinfectants for use against Foot and Mouth Disease and/or Swine Vesicular Disease relate to effectiveness when applied to a clean surface.

(a) thoroughly washed or sprayed with an approved disinfectant

(b) thoroughly cleansed, ensuring the dung, litter, etc. is removed and disposed of so there is no risk of contact with livestock; and

(c) (the clean area) washed or sprayed with an approved disinfectant used at the approved dilution rate.